



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

victim, which was over three inches longer than the "rattler," had been swallowed head foremost, and, because of its greater length lay in lateral folds in the stomach of the *Sistrurus*. As a result the latter snake was much distorted in outline. Due to handling and a consequent nervous reaction the Rattlesnake later disgorged its lengthy meal.

CLARENCE R. HALTER,
New York, N. Y.

RING-NECK SNAKE AT ORIENT, LONG ISLAND

In COPEIA, April, 1915, issue, a list of Long Island snakes is given with two stations for the Ring-neck Snake: Port Jefferson and Yaphank.

On September 12, 1915, while plowing in a weedy corner between two plots of woodlands, I turned out a young Ring-neck (*Diadophis punctatus*). It was five inches in length, of an uniform dark-gray above, the light band very prominent.

A small boy following in the furrow, who first called my attention to the specimen, reported that there were two of the same species; one quickly escaping into the thick bramble at the wood-margin, that one being much larger, as much as ten inches in length.

These were evidently hiding underneath loose turfts of sod. Much rubbish, boards, logs, stones and odds and ends are piled in the edge of the woods there. Although many searches have been made, no signs of others have appeared.

It is interesting to record a third station on Long Island for this secretive little reptile.

ROY LATHAM,
Orient, N. Y.

LOS ANGELES SALAMANDERS

In a branch canyon of Topango Canyon, near Santa Monica, California, is a small pool from which